Progress in Chemical Constituents and Pharmacological Effects of *Citrus medica* L. var. *sarcodactylis* Swingle

Shuyu Xu, Yin Qu, Xinyu Liu, Yao Li, Jimin Liu, and Hezhong Jiang*

School of Life Science and Engineering, Southwest Jiaotong University, Chengdu, Sichuan 611756, China

Email: jianghz10@swjtu.edu.cn (H. J.)

Abstract *Citrus medica* L. var. *sarcodactylis* Swingle is known as Foshou fruit that can be used as Chinese medicinal materials, and it has a health and longevity effect. In recent years, investigations about the chemical compounds of *Citrus medica* L. var. *sarcodactylis* Swingle have shown that its main compounds include flavonoids, essential oils, coumarins, and polysaccharides, with anti-inflammatory, antimicrobial, antioxidant activities, an effect on the autonomic nervous system, and other pharmacological effects. In this paper, the chemical compositions and pharmacological effects of *Citrus medica* L. var. *sarcodactylis* Swingle are summarized.

Keywords *Citrus medica* L. var. *sarcodactylis* Swingle, chemical constituents, pharmacological effects

Introduction

Citrus medica L. var. *sarcodactylis* Swingle (Rutaceae), one of the variants of *Citrus medica* L. var. *medica*, is an evergreen shrub or small tree and widely cultivated in Zhejiang, Fujian, Guangdong, Guangxi, Yunnan and Sichuan Province of China. *Citrus medica* L. var. *sarcodactylis* Swingle has the function of protecting liver, stomach and pancreas, and can treat hypertension and respiratory tract infection.^[1] In addition, it is traditionally used for the treatment of hypertension, respiratory tract infections and tracheitis in China.^[2] Due to the above effects, *Citrus medica* L. var. *sarcodactylis* Swingle has gradually attracted widespread attention in recent years. Herein, we summarize the chemical constituents and pharmacological effects of *Citrus medica* L. var. *sarcodactylis* Swingle.

Chemical Constituents of *Citrus medica* L. var. *sarcodactylis* Swingle

The chemical compositions of *Citrus medica* L. var. *sarcodactylis* Swingle are reported mainly including phenolic components, terpenoids, essential oil, and other components.^[3]

Phenolic constituent

The phenolic components of *Citrus medica* L. var. *sarcodactylis* Swingle are mainly flavonoids, coumarins, and other phenolic components (Figure 1).

The flavonoids are the main components in *Citrus medica* L. var. *sarcodactylis* Swingle. In 1984, He and Ling isolated two flavonoids from *Citrus medica* L. var. *sarcodactylis* Swingle, named 3,5,6-trihydroxy-4',7-dimethoxyflavone (1) and 3,5,6-trihydroxy-3',4',7'-trimethoxyflavone (2).^[4] Feng *et al.* isolated diosmetin (3), and diosmin (4) from the ethanol extract of *Citrus medica* L. var. *sarcodactylis* Swingle in 2004.^[5] In 2010, Chan *et al.* used a combination of traditional chromatography techniques isolated atalantoflavon (5), lonchocarpol A (6), and C-glycosylflavone-vitexin (7).^[6]

Coumarins are also important phenolic compounds of *Citrus medica* L. var. *sarcodactylis* Swingle. Gao *et al.* isolated 5,7-dimethoxycoumarin (8) from *Citrus medica* L. var. *sarcodactylis* Swingle in 2002.^[7] In 2010, Chan *et al.* purified a new coumarin named citrumedin-B (9) and xanthyletin (10),

Figure 1 The structures of phenolic components isolated from *Citrus medica* L. var. *sarcodactylis* Swingle.

nordentatin (11), 5,7-dimethoxycoumarin (12).^[6] In 2017, Chan *et al.* found 6,7-dimethoxycoumarin (13) from *Citrus medica* L. var. *sarcodactylis* Swingle. They proved that it has a significant inhibitory effect on superoxide anion formation.^[8]

In 2004, Feng *et al.* isolated aviprin (14), ferulic acid (15), vanillic acid (16), and protocatechuic acid (17) from the ethanol extract of *Citrus medica* L. var. *sarcodactylis* Swingle.^[5]

Terpenoids

The terpenoids in *Citrus medica* L. var. *sarcodactylis* Swingle are mainly triterpenoids especially tetranortriterpenoids, and sesquiterpenoids.

In 2010, Chan *et al.* purified thirty known compounds from *Citrus medica* L. var. *sarcodactylis* Swingle by using a combination of traditional chromatography techniques, including 1-(10 $\rightarrow$ 19)abeo-7 α -acetoxo-10 β -hydroxyisobacnoic acid-3,10-lactone (18), nomilin (19), limonin (20), lupeol

Figure 2 The structures of terpenoids isolated from *Citrus medica* L. var. *sarcodactylis* Swingle.

(21), limononic acid (22), citrussin (23) (Figure 2).^[6]

Then in 2017, Chan *et al.* found a new sesquiterpene named citrussin-C (24) by investigation of the chemical constituents from the fruits of *Citrus medica* L. var. *sarcodactylis* Swingle.^[8]

Essential oil

Essential oil is an important part of the study of *Citrus medica* L. var. *sarcodactylis* Swingle. Kim *et al.* studied the composition of essential oil of *Citrus medica* L. var. *sarcodactylis* Swingle by GC-MS, and identified fifteen components in 2013.^[9] According to their study, the main components of the essential oil of *Citrus medica* L. var. *sarcodactylis* Swingle are limonene (25) and γ -Terpinene (26) (Figure 3). Mitropoulou *et al.* used SPME GC-MS to analyze the main compounds of the essential oil of *C. medica* peels and assayed the antimicrobial properties in 2017.^[10] In 2018, Guo *et al.* compared chemical composition, antimicrobial and antioxidant activity of essential oil from *Citrus medica* L. var. *sarcodactylis* Swingle.^[11] They used GC-MS analysis for identifying more than 200 chemical components from the essential oil of *Citrus medica* L. var. *sarcodactylis* Swingle, which contained *D*-limonene (27), α -pinene (28), myrcene (29), ocimene (30), and linalool (31).

Figure 3 The structures of essential oil from *Citrus medica* L. var. *sarcodactylis* Swingle.

Other compounds

In addition to the above chemical components, Gao *et al.* isolated $\Delta^{5,22}$ -stigmaterol (32) and palmitic acid (33) from the ethyl acetate part of its ethanol extract in 2002. It is the first time $\Delta^{5,22}$ -stigmaterol was isolated from *Citrus medica* L. var. *sarcodactylis* Swingle.^[7] In 2004, Feng *et al.* isolated a new polysaccharide 5-methoxyfurfural and Kojic acid (34) from the ethanol extract of *Citrus medica* L. var. *sarcodactylis* Swingle.^[5]

Pharmacological Activities

Citrus medica L. var. *sarcodactylis* Swingle is a traditional Chinese medicinal plant with a variety of pharmacological activities such as antimicrobial, anti-inflammatory, antioxidative effect, and so on.^[12]

Antimicrobial activity

Theanphong *et al.* tested antimicrobial activity of essential oil with agar diffusion method and found its significant antimicrobial activity against *S. aureus* and *B. subtilis* with the

same minimum inhibitory concentration of 2500 ppm.^[13] Li *et al.* studied antimicrobial effect of *Citrus medica* L. var. *sarcodactylis* Swingle essential oil and its mechanism against food-borne bacteria. Then, they found that the essential oil showed a significant reduction effect on the growth rate of bacteria, and led to lysis of the cell wall, intracellular ingredient leakage, cell death eventually.^[14]

Anti-inflammatory activity

Kim *et al.* showed that the essential oil of *Citrus medica* L. var. *sarcodactylis* Swingle inhibited LPS-stimulated inflammation by blocking the NF- κ B, JNK, and ERK pathways in macrophages, and proved essential oil possessed anti-inflammatory properties.^[9]

Antioxidative activity

Zhu *et al.* found that the bergamot polysaccharide of *Citrus medica* L. var. *sarcodactylis* Swingle could efficiently remove O_2^- , HO^- and protect DNA chains significantly from being damaged by free radicals.^[15]

Effect on the autonomic nervous system

Sayorwan *et al.* investigated the *Citrus medica* L. var. *sarcodactylis* Swingle peel oil effect on the autonomic nervous systems and mood responses after inhaling.^[16]

Other pharmacological activities

Peng *et al.* proved that *Citrus medica* L. var. *sarcodactylis* Swingle fruit can promote insulin secretion by kinetic analysis on the hypoglycemic patterns of the intraperitoneal glucose tolerance (IPGTT) and the insulin-glucose tolerance tests (IGTT).^[1]

Conclusions and Perspectives

As one traditional Chinese medicine, *Citrus medica* L. var. *sarcodactylis* Swingle is distributed widely and it has a long medicinal history. The chemical constituents of *Citrus medica* L. var. *sarcodactylis* Swingle include polysaccharide, flavonoids, essential oils, coumarins, sesquiterpene, and other compounds. Pharmacodynamic studies show that it has a variety of pharmacological effects. At present, the research on *Citrus medica* L. var. *sarcodactylis* Swingle is mainly focused on the components and functions of essential oil, its other components and bioactivities need to be further studied.

Acknowledgement

This work was co-supported by the research project of the Fund of Science and Technology Agency of Chengdu (No. 2019-YF09-00049SN), the Administration of Traditional Chinese Medicine of Sichuan (No. 2017PC004), the 13th Personalized Experimental Project (Nos. GX201913085 and GX201913100), and the Student Research Training Program (SRTP, 2019156).

Author Contributions

Following are the details of the contributions made by each of the authors for the manuscript: Hezhong Jiang and Shuyu Xu performed the data analyses and wrote the manuscript; Yin Qu and Xinyu Liu contributed to the conception of the study; Yao Li and Jimin Liu contributed to analysis and manuscript preparation.

Conflict of Interest

The authors declare no conflict of interest.

Copyright © 2020 Shuyu Xu, Yin Qu, Xinyu Liu, Yao Li, Jimin Liu, and Hezhong Jiang. This article is an open access article distributed under

the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

References

- [1] Peng, C. H.; Ker, Y. B.; Weng, C. F. Insulin Secretagogue Bioactivity of Finger Citron Fruit (*Citrus medica* L. var. *Sarcodactylis* Hort, Rutaceae). *J. Agric. Food. Chem.* **2009**, *57*, 8812–8819.
- [2] He, Z.; Liang, F.; Zhang, Y. Water-soluble polysaccharides from finger citron fruits (*Citrus medica* L. var. *sarcodactylis*). *Carbohydr. Res.* **2014**, *388*, 100–104.
- [3] Nogata, Y.; Sakamoto, K.; Shiratsuchi, H. Flavonoid Composition of Fruit Tissues of Citrus Species. *Biosci. Biotechnol. Biochem.* **2006**, *70*, 178–192.
- [4] He, H. Y.; Ling, L. Q. Chemical studies on a Chinese traditional drug fingered citron (*Citrus medica* L. var. *sarcodactylis* (Noot.) Swingle). *Acta Pharm. Sinica* **1985**, *6*, 433–435.
- [5] Yin, F.; Lou, F. Studies on the Constituents of *Citrus medica* L. var. *Sarcodactylis* (Noot.) Swingle. *Chin. Pharm. J.* **2004**, *39*, 24–26.
- [6] Chan, Y. Y.; Li, C. H.; Shen, Y. C. Anti-inflammatory Principles from the Stem and Root Barks of *Citrus medica*. *Chem. Pharm. Bull.* **2010**, *58*, 61–65.
- [7] Gao, Y. H.; Xu, H. H.; Diao, Y. M. Studies on Chemical Components of *Citrus medica* L. var. *Sarcodactylis* Swingle. *Chin. Med. Mater.* **2002**, *13*, 315–316.
- [8] Chan, Y. Y.; Hwang, T. L.; Kuo, P. Constituents of the Fruits of *Citrus medica* L. var. *sarcodactylis* and the Effect of 6,7-Dimethoxy-coumarin on Superoxide Anion Formation and Elastase Release. *C. Molecules* **2017**, *22*, 1454–1463.
- [9] Kim, K. N.; Ko, Y. J.; Yang, H. M. Anti-inflammatory effect of essential oil and its constituents from fingered citron (*Citrus medica* L. var. *sarcodactylis*) through blocking JNK, ERK and NF- κ B signaling pathways in LPS-activated RAW 264.7 cells. *Food Chem. Toxicol.* **2013**, *57*, 126–131.
- [10] Mitropoulou, G.; Fitsiou, E.; Spyridopoulou, K. Citrus medica essential oil exhibits significant antimicrobial and antiproliferative activity. *Lwt-Food Sci. Technol.* **2017**, *84*, 344–352.
- [11] Guo, J. J.; Gao, Z. P.; Xia, J. L. Comparative analysis of chemical composition, antimicrobial and antioxidant activity of citrus essential oils from the main cultivated varieties in China. *Lwt-Food Sci. Technol.* **2018**, *97*, 825–839.
- [12] Sakurada, T.; Kuwahata, H.; Katsuyama, S. Intraplantar injection of bergamot essential oil into the mouse hindpaw: effects on capsaicin-induced nociceptive behaviors. *Int. Rev. Neurobiol.* **2009**, *85*, 237–248.
- [13] Theanphong, O.; Songsak, T.; Mingvanish, W. Chemical Composition and Antimicrobial Activity of the Essential Oil from *Citrus medica* L. var. *sarcodactylis* (Sieber) Swingle Leaf. *Mahidol Univ. J. Pharm. Sci.* **2008**, *35*, 57–61.
- [14] Li, Z. H.; Cai, M.; Liu, Y. S. Antibacterial Activity and Mechanisms of Essential Oil from *Citrus medica* L. var. *sarcodactylis*. *Molecules* **2019**, *24*, 1577–1587.
- [15] Zhu, X. Y.; Wu, J. M.; Jia, Z. S. Composition and Antioxidative Activity of Polysaccharide from Bergamot. *Chem. J. Chin. Univ.* **2005**, *26*, 1264–1267.
- [16] Sayorwan, W.; Rumruay, V. Chemical compositions of fingered citron peel (*Citrus medica* L. var. *sarcodactylis*) and its effect on the autonomic nervous system. *J. Health Res.* **2017**, *31*, 307–313.

Received December 23, 2019

Accepted January 17, 2020